

An AP-Structure With Finslerian Flavor II: Torsion, Curvature and Other Objects

M.I.Wanas^{1,3,4} and Mona M. Kamal^{2,3}

Abstract

An absolute parallelism (AP-) space having Finslerian properties is called FAP-space. This FAP-structure is more wider than both conventional AP and Finsler structures. In the present work, more geometric objects as curvature and torsion tensors are derived in the context of this structure. Also second order tensors, usually needed for physical applications, are derived and studied. Furthermore, the anti-curvature and the W-tensor are defined for the FAP-structure. Relations between Riemannian, AP, Finsler and FAP structures are given. These relations facilitate comparison between results of applications carried out in the framework of these structures. We hope that the use of the FAP-structure, in applications may throw some light on some of the problems facing geometric field theories.

Keywords: AP-Geometry, Finsler Geometry, W-tensor, Anti-curvature tensor.

PACS: 02.40.Hw

1 Introduction and Motivations

Recently, a lot of attention has been paid to Finsler geometry for applications (cf. Ref. 1, 2, 3). On the other hand, teleparallel geometry (absolute parallelism (AP-) geometry) has gained many authors (cf. Ref. 4, 5, 6) for the purpose of physical applications. The interest in such geometries has been grown for the goal of solving problems of gravity theories, recently discussed (cf. Ref. 7, 8, 9). The aim has been to construct field theories, including general relativity (GR), in more wider geometries than the Riemannian one.

In this direction, one of the authors (MIW) has attempted to find a type of geometry with both AP and Finsler properties (FAP-structure). The geometric structure proposed [10] is more wider than both AP and Finsler geometries. The present work is the second part of this work, in which torsion, curvature and other objects are derived in the FAP. The motivation of this work is given, in some details, in the first part [10].

The paper is organized in the following manner. In the next Section, we give a brief review of the results obtained in the first part. Second order tensors and scalars, that may be important for physical applications, are given in Section 3. In Section 4, torsion, curvature and anti-curvature tensors, of the FAP-structure, are derived. In Section 5, the W-tensor is derived for the present structure. The paper is discussed and concluded in Section 6. In what follows we are going to characterize geometric objects, in the FAP-structure, by putting a hat on the same notations used in AP-geometry.

¹Astronomy Department, Faculty of Science, Cairo University, Giza, Egypt.

E-mail:mwanas@cu.edu.eg; wanas@frcu.eun.eg

²Mathematics Department, Faculty of Girls, Ain Shams University, Cairo, Egypt.

E-mail:monamkamal@eun.eg

³Egyptian Relativity Group (ERG) URL:<http://www.erg.eg.net>

⁴Center for Theoretical Physics (CTP) at the British University in Egypt (BUE).

2 Summary of The Results of the First Part

In first part [10] the following results have been obtained:

Definition: An Absolute Parallelism space with Finslerian (FAP) properties (M, L_i) is an n -dimensional differentiable manifold M equipped with a set of n-Lagrangian functions $L_i \equiv L_i(x, y)$ having the following properties:

1. $L_i(x, y)$ is C^∞ on $\tau M (= TM \setminus \{0\})$.
2. $L_i(x, y) > 0, y \in \tau M, y = \dot{x} (= \frac{dx}{dt})$.
3. $L_i(x, y)$ is positively homogenous of degree one ($P - h(1)$).
4. The vector fields ⁴

$$\hat{\lambda}_\mu^\mu(x, y) \stackrel{\text{def.}}{=} \frac{\partial L_i}{\partial y^\mu} = L_{i;\mu}, \quad (1)$$

are assumed to be linearly independent and globally defined on M , then we can define the conjugate vector fields $\hat{\lambda}_i^\mu$ such that,

$$\hat{\lambda}_i^\mu \hat{\lambda}_\nu^\mu = \delta_\nu^\mu, \quad (2)$$

$$\hat{\lambda}_i^\mu \hat{\lambda}_j^\mu = \delta_{ij}. \quad (3)$$

These vector fields are the building blocks of the FAP-Structure. The following object,

$$N_{\cdot\alpha}^\nu \stackrel{\text{def.}}{=} y^\mu \hat{\lambda}_i^\nu \hat{\lambda}_{\mu,\alpha}, \quad (4)$$

is found to transform as a *non-linear connection*. The following second order tensor,

$$C_{i\mu\alpha} \stackrel{\text{def.}}{=} \frac{\partial \hat{\lambda}_{i\mu}}{\partial y^\alpha} = L_{i;\mu\alpha}, \quad (5)$$

plays an important role in the FAP-Structure. It is type (0,2) symmetric tensor and P-h(-1). The following theorem has been proved.

Theorem 1. "A necessary and sufficient condition for an FAP-space to be an AP-space is that $C_{\alpha\beta}$ vanishes identically".

It is shown that, if

$$\hat{C}_{\cdot\alpha\beta}^\mu \stackrel{\text{def.}}{=} \hat{\lambda}_i^\mu C_{\alpha\beta}, \quad (6)$$

then,

$$C_{\alpha\beta} = \hat{\lambda}_i^\mu \hat{C}_{\cdot\alpha\beta}^\mu. \quad (7)$$

⁴In what follows the colon (:) is used for differentiation w.r.t. y , while comma (,) is used for differentiation w.r.t. x , $\mu (= 1, 2, \dots, n)$ represents the vector component and $i (= 1, 2, \dots, n)$ is the vector number.

Using the non-linear connection (4), we can define the following operator

$$\delta_\mu \stackrel{\text{def.}}{=} \partial_\mu - N_{\cdot\mu}^\alpha \frac{\partial}{\partial y^\alpha}. \quad (8)$$

This operator can be written as $(\hat{\cdot})$, if it is used as an infix operator.

The following geometric objects have been shown to transform as linear connections in the FAP-space:

1. *Cartan-Like* **Linear Connection**,

$$\hat{\Gamma}_{\cdot\alpha\beta}^\mu \stackrel{\text{def.}}{=} \hat{\lambda}_i^\mu \hat{\lambda}_{\alpha\hat{\cdot}\beta}^\mu. \quad (9)$$

Since (9) is non-symmetric, we can define the dual linear connection,

$$\tilde{\hat{\Gamma}}_{\cdot\alpha\beta}^\mu \stackrel{\text{def.}}{=} \hat{\Gamma}_{\cdot\beta\alpha}^\mu, \quad (10)$$

and also the symmetric linear connection⁵,

$$\hat{\Gamma}_{\cdot(\alpha\beta)}^\mu \stackrel{\text{def.}}{=} \frac{1}{2}(\hat{\Gamma}_{\cdot\alpha\beta}^\mu + \hat{\Gamma}_{\cdot\beta\alpha}^\mu). \quad (11)$$

2. *Bervald-Like* **Linear Connection**,

$$\hat{B}_{\cdot\alpha\beta}^\mu \stackrel{\text{def.}}{=} N_{\alpha\hat{\cdot}\beta}^\mu = \frac{\partial}{\partial y^\beta} (y^\nu \hat{\lambda}_i^\mu \hat{\lambda}_{\nu\hat{\cdot}\alpha}^\mu). \quad (12)$$

Connections (9), (10) and (12) are non-symmetric in their lower indices (α, β) . Consequently, they possess torsion, respectively.

Using the objects (9), (10), (11) and (6) one can define the following covariant derivatives, respectively:

If A_μ is an arbitrary vector, then its horizontal (H-) positive derivative is defined by

$$A_{\alpha||\beta} \stackrel{\text{def.}}{=} A_{\alpha\hat{\cdot}\beta} - A_\mu \hat{\Gamma}_{\cdot\alpha\beta}^\mu. \quad (13)$$

Also, we can define the following H-negative derivative using the dual connection, as follows

$$A_{\alpha||\beta} \stackrel{\text{def.}}{=} A_{\alpha\hat{\cdot}\beta} - A_\mu \tilde{\hat{\Gamma}}_{\cdot\alpha\beta}^\mu. \quad (14)$$

In addition we can define a third H-neutral derivative using the symmetric part of the *Cartan like* linear connection, i.e.

$$A_{\alpha||\beta} \stackrel{\text{def.}}{=} A_{\alpha\hat{\cdot}\beta} - A_\mu \hat{\Gamma}_{\cdot(\alpha\beta)}^\mu. \quad (15)$$

⁵The parentheses () stands for the symmetric part while the brackets [] stands for the skew part, for any geometric object, w.r.t. the indices included.

The vertical (V-) derivative of the vector A_μ , is defined by

$$A_\mu|_\nu \stackrel{\text{def.}}{=} A_{\mu;\nu} - A_\alpha \hat{C}^\alpha_{\cdot\mu\nu}. \quad (16)$$

Since $\hat{C}^\alpha_{\cdot\mu\nu}$ is symmetric in its two lower indices (μ, ν) , then we have only one type of derivative.

It has been shown that,

$$\hat{\lambda}_\mu|_\nu \equiv 0, \quad (17)$$

$$\hat{\lambda}_{\mu||\nu} \equiv 0. \quad (18)$$

Relations (17) and (18) show clearly that $\hat{\lambda}_\mu$ are parallel displaced along certain paths characterized by the geometric objects (6) and (9), respectively.

The second order symmetric tensors,

$$\hat{g}_{\mu\nu} \stackrel{\text{def.}}{=} \hat{\lambda}_\mu \hat{\lambda}_\nu, \quad (19)$$

$$\hat{g}^{\alpha\beta} \stackrel{\text{def.}}{=} \hat{\lambda}^\alpha \hat{\lambda}^\beta, \quad (20)$$

can be used to play the role of the metric tensor (and its conjugate) of Finsler space associated with the FAP-space, under certain condition [10]. It is clear from the definition (19) and the identities (18) that

$$\hat{g}_{\mu\nu||\sigma} \equiv 0, \quad (21)$$

which means that the linear connection (9) is a metric one. Also, it is shown that

$$\hat{g}_{\mu\nu}|_\sigma \equiv 0. \quad (22)$$

Relations (21), (22) means simply that the operation of raising and lowering tensor indices commutes with the operations of H-positive and V- differentiation.

Further Relations

If we define the tensor,

$$\hat{C}^{\star \mu}_{\cdot \alpha} \stackrel{\text{def.}}{=} \hat{\lambda}^\mu_{\cdot \alpha}, \quad (23)$$

then we can write,

$$\hat{C}^{\star \mu}_{\cdot \nu\alpha} \stackrel{\text{def.}}{=} \hat{\lambda}^\mu_{\cdot \nu} \hat{\lambda}_\alpha, \quad (24)$$

which in virtue of (2) and definition (6) can be written as,

$$\hat{C}^{\star \mu}_{\cdot \nu\alpha} = - \hat{C}^\mu_{\cdot \nu\alpha}.$$

Now, since $\hat{C}^\mu_{\cdot \nu\alpha}$ is symmetric w.r.t. its two lower indices, then we get,

$$\hat{C}^{\star \mu}_{\cdot [\nu\alpha]} = 0,$$

which gives immediately the relation,

$$\hat{\lambda}_{i\cdot}^{\mu}{}_{:\alpha} \hat{\lambda}_i^{\nu} = \hat{\lambda}_{i\cdot}^{\mu}{}_{:\nu} \hat{\lambda}_i^{\alpha}. \quad (25)$$

This is an important relation characterizing the FAP-structure.

The canonical linear connection (9) can be expressed in the form

$$\hat{\Gamma}_{\cdot\mu\nu}^{\alpha} \stackrel{\text{def.}}{=} \left\{ \begin{matrix} \hat{\alpha} \\ \mu\nu \end{matrix} \right\} + \hat{\gamma}_{\cdot\mu\nu}^{\alpha}, \quad (26)$$

where $\left\{ \begin{matrix} \hat{\alpha} \\ \mu\nu \end{matrix} \right\}$ is the linear metric connection defined by,

$$\left\{ \begin{matrix} \hat{\alpha} \\ \mu\nu \end{matrix} \right\} \stackrel{\text{def.}}{=} \frac{1}{2} \hat{g}^{\alpha\beta} (\hat{g}_{\beta\mu;\nu} + \hat{g}_{\beta\nu;\mu} - \hat{g}_{\mu\nu;\beta}),$$

and $\hat{\gamma}_{\cdot\mu\nu}^{\alpha}$ is a third order tensor defined by,

$$\hat{\gamma}_{\cdot\mu\nu}^{\alpha} \stackrel{\text{def.}}{=} \hat{\lambda}_i^{\alpha} \hat{\lambda}_{\mu;\nu}^{\hat{i}}, \quad (27)$$

the infix operator $(\hat{;})$ stands for the covariant differential operator using $\left\{ \begin{matrix} \hat{\alpha} \\ \mu\nu \end{matrix} \right\}$ i.e.

$$\hat{\lambda}_{\mu;\nu}^{\hat{i}} \stackrel{\text{def.}}{=} \hat{\lambda}_i^{\hat{i}\mu;\nu} - \hat{\lambda}_i^{\epsilon} \left\{ \begin{matrix} \hat{\epsilon} \\ \mu\nu \end{matrix} \right\} \quad (28)$$

For the FAP-structure we have a set of, at least, four d-connections: $D = (\left\{ \begin{matrix} \hat{\alpha} \\ \mu\nu \end{matrix} \right\}, N_{\mu}^{\alpha}, \hat{C}_{\cdot\mu\nu}^{\alpha})$, $\hat{D} = (\hat{\Gamma}_{\cdot\mu\nu}^{\alpha}, N_{\mu}^{\alpha}, \hat{C}_{\cdot\mu\nu}^{\alpha})$, $\tilde{D} = (\tilde{\Gamma}_{\cdot\mu\nu}^{\alpha}, N_{\mu}^{\alpha}, \hat{C}_{\cdot\mu\nu}^{\alpha})$ and $\overline{D} = (\hat{\Gamma}_{\cdot(\mu\nu)}^{\alpha}, N_{\mu}^{\alpha}, \hat{C}_{\cdot\mu\nu}^{\alpha})$.

3 Tensors of Different Orders

In geometric field theories, different geometric objects play important roles in representing physical quantities, in the context of the geometrization philosophy. To describe different physical fields, one may need scalars, vectors and /or second order tensors. On the other hand, third order objects (connections or tensors) are necessary to describe trajectories of test particles in different physical fields. For these reasons, we are going to give definitions of such objects in the FAP-Structure. This is done in order to make this structure ready for applications. Of course, in what follows we cannot cover all such objects, but only those that we expect, from previous experience, to play a role in physical applications.

3.1 Third Order Tensors and Vectors

Now, we are going to give definitions for some important third order tensors and their contractions. The torsion of the linear connection (9) is given by,

$$\hat{\Lambda}_{\cdot\mu\nu}^{\alpha} \stackrel{\text{def.}}{=} \hat{\Gamma}_{\cdot\mu\nu}^{\alpha} - \hat{\Gamma}_{\cdot\nu\mu}^{\alpha} = \hat{\gamma}_{\cdot\mu\nu}^{\alpha} - \hat{\gamma}_{\cdot\nu\mu}^{\alpha}, \quad (29)$$

which gives by contraction the vector,

$$\hat{c}_\alpha \stackrel{\text{def.}}{=} \hat{\Lambda}_{\alpha\mu}^\mu = \hat{\gamma}_{\alpha\mu}^\mu. \quad (30)$$

Also, the torsion of the *Bervard-like* connection (12) is defined by,

$$\hat{T}_{\mu\nu}^\alpha \stackrel{\text{def.}}{=} \hat{B}_{\mu\nu}^\alpha - \hat{B}_{\nu\mu}^\alpha, \quad (31)$$

which gives the vector,

$$\hat{T}_\mu \stackrel{\text{def.}}{=} \hat{T}_{\mu\alpha}^\alpha. \quad (32)$$

Now, using the contortion (27), we can define the third order tensors,

$$\hat{\Delta}_{\mu\nu}^\alpha \stackrel{\text{def.}}{=} \hat{\gamma}_{\mu\nu}^\alpha + \hat{\gamma}_{\nu\mu}^\alpha, \quad (33)$$

$$\hat{\gamma}_{\mu\nu}^{\alpha} \stackrel{\text{def.}}{=} \hat{g}^{\epsilon\alpha} \hat{g}_{\mu\beta} \hat{\gamma}_{\nu\epsilon}^\beta \quad (34)$$

Also, using the non-linear connection (4) and the operator (8), we can define the object,

$$N_{\mu\nu}^\alpha \stackrel{\text{def.}}{=} N_{\mu\hat{\nu}}^\alpha - N_{\hat{\nu}\mu}^\alpha. \quad (35)$$

This object can be shown (see Sec. 4) to be tensor of type (1,2). Its contraction gives the vector,

$$N_\mu \stackrel{\text{def.}}{=} N_{\mu\alpha}^\alpha. \quad (36)$$

The contraction of the tensor (6) will give the vector,

$$\hat{V}_\alpha \stackrel{\text{def.}}{=} \hat{C}_{\alpha\mu}^\mu \quad (37)$$

3.2 Second Order Tensors and Scalars

In the AP-geometry, a set of second order tensors has been defined [11]. Such tensors have been used to classify [12], physically, different AP-structures (with different symmetries) used for applications. This classification scheme is a covariant one [13]. It gives the capabilities of certain AP-structures to represent physical situations. Also, some physical meaning have been attributed to these tensors. Such tensors are obtained using the torsion, contortion tensors of the linear connection and their contraction. So, it would be interesting to define the corresponding second order tensors in the context of the FAP-structure. This would facilitate comparison with the physical results obtained from Ap-geometry, when using FAP-geometry for physical applications.

The following Table gives a summary of second order tensors defined in the context of the FAP-geometry. As stated above, we use the same notation of the AP-geometry, with a hat characterizing the FAP-geometry. The condition for removing this hat is given by Theorem 1. Table 1 is similar to that of the AP-structure [11].

Table 1: Second Order World Tensors

Skew Tensors	Symmetric Tensors
$\hat{\xi}_{\mu\nu} \stackrel{\text{def.}}{=} \hat{\gamma}_{\mu\nu, \epsilon}^\epsilon$	
$\hat{\zeta}_{\mu\nu} \stackrel{\text{def.}}{=} \hat{c}_\alpha \hat{\gamma}_{\mu\nu}^\alpha$	
$\hat{\eta}_{\mu\nu} \stackrel{\text{def.}}{=} \hat{c}_\alpha \hat{\Lambda}_{\mu\nu}^\alpha$	$\hat{\phi}_{\mu\nu} \stackrel{\text{def.}}{=} \hat{c}_\alpha \hat{\Delta}_{\mu\nu}^\alpha$
$\hat{\chi}_{\mu\nu} \stackrel{\text{def.}}{=} \hat{\Lambda}_{\mu\nu, \epsilon}^\epsilon$	$\hat{\psi}_{\mu\nu} \stackrel{\text{def.}}{=} \hat{\Delta}_{\mu\nu, \epsilon}^\epsilon$
$\hat{\varepsilon}_{\mu\nu} \stackrel{\text{def.}}{=} \hat{c}_{\mu \nu} - \hat{c}_{\nu \mu}$	$\hat{\theta}_{\mu\nu} \stackrel{\text{def.}}{=} \hat{c}_{\mu \nu} + \hat{c}_{\nu \mu}$
$\hat{\kappa}_{\mu\nu} \stackrel{\text{def.}}{=} \hat{\gamma}_{\mu\epsilon}^\alpha \hat{\gamma}_{\alpha\nu}^\epsilon - \hat{\gamma}_{\nu\epsilon}^\alpha \hat{\gamma}_{\alpha\mu}^\epsilon$	$\hat{\omega}_{\mu\nu} \stackrel{\text{def.}}{=} \hat{\gamma}_{\mu\epsilon}^\alpha \hat{\gamma}_{\alpha\nu}^\epsilon + \hat{\gamma}_{\nu\epsilon}^\alpha \hat{\gamma}_{\alpha\mu}^\epsilon$
	$\hat{\omega}_{\mu\nu} \stackrel{\text{def.}}{=} \hat{\gamma}_{\mu\alpha}^\epsilon \hat{\gamma}_{\nu\epsilon}^\alpha$
	$\hat{\sigma}_{\mu\nu} \stackrel{\text{def.}}{=} \hat{\gamma}_{\alpha\mu}^\epsilon \hat{\gamma}_{\epsilon\nu}^\alpha$
	$\hat{\alpha}_{\mu\nu} \stackrel{\text{def.}}{=} \hat{c}_\mu \hat{c}_\nu$

where $\hat{\gamma}_{\mu\nu,||\epsilon}^\epsilon \stackrel{\text{def.}}{=} \hat{\gamma}_{\mu\nu,++||\epsilon}^\epsilon$ and so on.

We can obtain another set of second order tensors, resulting from the combination of (6), (29), (31), (35) and their contractions, (e.g. skew-tensors, $(N_\alpha \hat{\gamma}_{\mu\nu}^\alpha, N_\alpha \hat{\Lambda}_{\mu\nu}^\alpha, \hat{c}_\alpha \hat{T}_{\mu\nu}^\alpha, N_\alpha N_{\mu\nu}^\alpha, \dots)$, and symmetric-tensors, $(\hat{V}_\alpha \hat{V}_\beta, \hat{c}_\alpha \hat{C}_{\mu\nu}^\alpha, \hat{T}_\alpha \hat{C}_{\mu\nu}^\alpha, \hat{T}_\mu \hat{T}_\nu, \dots)$). The properties of such tensors will be studied in details, after application, depending on their importance for such applications.

From the tensors given in Table 1, we can define many scalars (e.g. $\hat{\phi} \stackrel{\text{def.}}{=} \hat{c}_\alpha \hat{\Delta}_{\mu\nu}^{\alpha\mu}$, $\hat{\psi} \stackrel{\text{def.}}{=} \hat{\Delta}_{\mu\nu,||\epsilon}^{\epsilon\mu}$, $\hat{\theta} \stackrel{\text{def.}}{=} 2 \hat{c}_{\mu||\mu}^\mu$, $\hat{\alpha} \stackrel{\text{def.}}{=} \hat{c}^\mu \hat{c}_\mu$, $\hat{\omega} \stackrel{\text{def.}}{=} \hat{\gamma}_{\mu\epsilon}^{\alpha\mu} \hat{\gamma}_{\alpha\mu}^\epsilon + \hat{\gamma}_{\mu\epsilon}^\alpha \hat{\gamma}_{\alpha\mu}^{\epsilon\mu}$, $\hat{\omega} \stackrel{\text{def.}}{=} \hat{\gamma}_{\mu\epsilon}^{\epsilon\mu} \hat{\gamma}_{\alpha\mu}^\alpha$ and $\hat{\sigma} \stackrel{\text{def.}}{=} \hat{\gamma}_{\alpha\mu}^\epsilon \hat{\gamma}_{\epsilon\mu}^{\alpha\mu}$).

Among these tensors, we have a set of second order traceless tensors (e.g. $(\hat{\phi}_{\mu\nu} + \hat{\alpha}_{\mu\nu})$, $(\hat{\psi}_{\mu\nu} + \hat{\theta}_{\mu\nu})$ and $(\hat{\omega}_{\mu\nu} + 2 \hat{\omega}_{\mu\nu})$) with possible linear combination. Such object may be of physical interest.

4 Curvature, Anti-Curvature and Torsion Tensors

It is well known that any differentiable manifold equipped with a d-connection, $F = (F_{\mu\nu}^\alpha, G_{\beta\gamma}^\alpha, C_{\mu\nu}^\alpha)$, implies the following commutation relations (cf. Ref. 14)

$$A_\alpha|_{\beta\sigma} - A_\alpha|_{\sigma\beta} = A_\mu R_{\alpha\beta\sigma}^\mu - A_\alpha|_\mu (G_{\beta\sigma}^\mu - G_{\sigma\beta}^\mu) - A_\alpha|_\mu \Lambda_{\beta\sigma}^\mu,$$

$$A_\alpha|_{\beta\sigma} - A_\alpha|_{\sigma\beta} = A_\mu S_{\alpha\beta\sigma}^\mu - A_\alpha|_\mu (C_{\beta\sigma}^\mu - C_{\sigma\beta}^\mu),$$

where A_μ is an arbitrary vector field, function of both x and y , $C_{\mu\nu}^\alpha$ is 3rd order tensor, $G_{\beta\gamma}^\alpha$ is a non-linear connection, $R_{\alpha\beta\sigma}^\mu$ is the curvature using the linear connection $F_{\mu\nu}^\alpha$ and $\Lambda_{\beta\sigma}^\mu$ is the torsion of this connection. Also, $|$ and $|$ denote differentiation using $F_{\mu\nu}^\alpha$ and $C_{\mu\nu}^\alpha$, respectively.

Now using distinguish connections, $\hat{D}$, $\tilde{D}$, $\overline{D}$ and D of the FAP-structure, we can write the following horizontal commutation relations, respectively,

$$A_{\alpha||\beta\sigma} - A_{\alpha||\sigma\beta} = A_{\mu}\hat{B}^{\mu}_{\alpha\beta\sigma} - A_{\alpha}N^{\epsilon}_{\beta\sigma} - A_{\alpha||\mu}\hat{\Lambda}^{\mu}_{\beta\sigma}, \quad (38)$$

$$A_{\alpha||\beta\sigma} - A_{\alpha||\sigma\beta} = A_{\mu}\tilde{B}^{\mu}_{\alpha\beta\sigma} - A_{\alpha}N^{\epsilon}_{\beta\sigma} - A_{\alpha||\mu}\hat{\Lambda}^{\mu}_{\beta\sigma}, \quad (39)$$

$$A_{\alpha||\beta\sigma} - A_{\alpha||\sigma\beta} = A_{\mu}\overline{B}^{\mu}_{\alpha\beta\sigma} - A_{\alpha}N^{\epsilon}_{\beta\sigma}, \quad (40)$$

$$A_{\alpha\hat{;}\beta\sigma} - A_{\alpha\hat{;}\sigma\beta} = A_{\mu}\hat{R}^{\mu}_{\alpha\beta\sigma} - A_{\alpha}N^{\epsilon}_{\beta\sigma}. \quad (41)$$

Also, using the vertical derivative, we can write,

$$A_{\alpha}|\beta\sigma - A_{\alpha}|\sigma\beta = A_{\mu}\hat{S}^{\mu}_{\alpha\beta\sigma}, \quad (42)$$

where,

$$\hat{B}^{\mu}_{\alpha\beta\sigma} \stackrel{\text{def.}}{=} \hat{\Gamma}^{\mu}_{\alpha\sigma\hat{;}\beta} - \hat{\Gamma}^{\mu}_{\alpha\beta\hat{;}\sigma} + \hat{\Gamma}^{\mu}_{\epsilon\beta}\hat{\Gamma}^{\epsilon}_{\alpha\sigma} - \hat{\Gamma}^{\mu}_{\epsilon\sigma}\hat{\Gamma}^{\epsilon}_{\alpha\beta} - \hat{C}^{\mu}_{\alpha\epsilon}N^{\epsilon}_{\beta\sigma}, \quad (43)$$

$$\tilde{B}^{\mu}_{\alpha\beta\sigma} \stackrel{\text{def.}}{=} \tilde{\Gamma}^{\mu}_{\alpha\sigma\hat{;}\beta} - \tilde{\Gamma}^{\mu}_{\alpha\beta\hat{;}\sigma} + \tilde{\Gamma}^{\mu}_{\epsilon\beta}\tilde{\Gamma}^{\epsilon}_{\alpha\sigma} - \tilde{\Gamma}^{\mu}_{\epsilon\sigma}\tilde{\Gamma}^{\epsilon}_{\alpha\beta} - \hat{C}^{\mu}_{\alpha\epsilon}N^{\epsilon}_{\beta\sigma}, \quad (44)$$

$$\overline{B}^{\mu}_{\alpha\beta\sigma} \stackrel{\text{def.}}{=} \hat{\Gamma}^{\mu}_{(\alpha\sigma)\hat{;}\beta} - \hat{\Gamma}^{\mu}_{(\alpha\beta)\hat{;}\sigma} + \hat{\Gamma}^{\mu}_{(\epsilon\beta)}\hat{\Gamma}^{\epsilon}_{(\alpha\sigma)} - \hat{\Gamma}^{\mu}_{(\epsilon\sigma)}\hat{\Gamma}^{\epsilon}_{(\alpha\beta)} - \hat{C}^{\mu}_{\alpha\epsilon}N^{\epsilon}_{\beta\sigma}, \quad (45)$$

$$\hat{R}^{\mu}_{\alpha\beta\sigma} \stackrel{\text{def.}}{=} \left\{ \begin{matrix} \hat{\mu} \\ \alpha\sigma \end{matrix} \right\}_{\hat{;}\beta} - \left\{ \begin{matrix} \hat{\mu} \\ \alpha\beta \end{matrix} \right\}_{\hat{;}\sigma} + \left\{ \begin{matrix} \hat{\mu} \\ \epsilon\beta \end{matrix} \right\} \left\{ \begin{matrix} \hat{\epsilon} \\ \alpha\sigma \end{matrix} \right\} - \left\{ \begin{matrix} \hat{\mu} \\ \epsilon\sigma \end{matrix} \right\} \left\{ \begin{matrix} \hat{\epsilon} \\ \alpha\beta \end{matrix} \right\} - \hat{C}^{\mu}_{\alpha\epsilon}N^{\epsilon}_{\beta\sigma}, \quad (46)$$

and

$$\hat{S}^{\mu}_{\alpha\beta\sigma} \stackrel{\text{def.}}{=} \hat{C}^{\mu}_{\alpha\sigma\hat{;}\beta} - \hat{C}^{\mu}_{\alpha\beta\hat{;}\sigma} + \hat{C}^{\mu}_{\epsilon\beta}\hat{C}^{\epsilon}_{\alpha\sigma} - \hat{C}^{\mu}_{\epsilon\sigma}\hat{C}^{\epsilon}_{\alpha\beta}. \quad (47)$$

It can be shown; applying the *Quotient Theorem* on the terms containing $N^{\alpha}_{\mu\nu}$, appearing in (38), (39), (40), (41) and defined by (35); that $N^{\alpha}_{\mu\nu}$ is a third order tensor, skew symmetric in its lower indices. The geometric quantities defined by (43), (44), (45), (46) and (47) are curvature tensors of the d-connections of the FAP-structure, respectively.

Theorem 2. "The horizontal curvature (43) vanishes identically" i.e. $\hat{B}^{\mu}_{\alpha\beta\sigma} \equiv 0$.
This theorem can be proved using definitions (9) and (18).

Theorem 3. "The vertical curvature (47) vanishes identically" i.e. $\hat{S}^{\mu}_{\alpha\beta\sigma} \equiv 0$.
This theorem can be easily proved using definition (6).

The curvature tensor $\hat{B}^{\mu}_{\alpha\beta\sigma}$ (43) can be expressed in terms of the contortion tensor (27) and curvature (46) i.e.

$$\hat{B}^{\mu}_{\alpha\beta\sigma} = \hat{R}^{\mu}_{\alpha\beta\sigma} + \hat{Q}^{\mu}_{\alpha\beta\sigma} \quad (48)$$

where,

$$\hat{Q}^{\mu}_{\alpha\beta\sigma} \stackrel{\text{def.}}{=} \hat{\gamma}^{\mu}_{\alpha\sigma||\beta} - \hat{\gamma}^{\mu}_{+\alpha\beta||\sigma} - \hat{\gamma}^{\nu}_{\alpha\sigma}\hat{\gamma}^{\mu}_{\nu\beta} + \hat{\gamma}^{\nu}_{\alpha\beta}\hat{\gamma}^{\mu}_{\nu\sigma}, \quad (49)$$

and $\hat{R}^{\mu}_{\alpha\beta\sigma}$ is defined by (46). Now from the definitions (49) and (46), it is clear that neither $\hat{Q}^{\mu}_{\alpha\beta\sigma}$ nor $\hat{R}^{\mu}_{\alpha\beta\sigma}$ vanishes, while their sum, given by (48) vanishes (Theorem 2). It is to be considered that, the curvature tensor $\hat{R}^{\mu}_{\alpha\beta\sigma}$ is made of the linear connection $\{\hat{\gamma}^{\mu}_{\alpha\beta}\}$ and the tensors $\hat{C}^{\mu}_{\alpha\beta}$ and $N^{\mu}_{\alpha\beta}$ while the tensor $\hat{Q}^{\mu}_{\alpha\beta\sigma}$ is purely made of the torsion (or the contortion) of the canonical connection. The addition of these two tensors gives rise to the vanishing of the total curvature of the space (Theorem 2). For the above mentioned comments, we call $\hat{Q}^{\mu}_{\alpha\beta\sigma}$ the "additive inverse of the curvature tensor" [15] or the "anti-curvature" tensor [16].

The following Table gives the possible contractions of the curvature tensors $\tilde{\hat{B}}^{\mu}_{\alpha\beta\sigma}$ (44), $\overline{\hat{B}}^{\mu}_{\alpha\beta\sigma}$ (45) and $\hat{R}^{\mu}_{\alpha\beta\sigma}$ (46) in terms of the second order tensors of Table 1. Note for example: $\tilde{\hat{K}}_{\beta\sigma} \stackrel{\text{def.}}{=} \tilde{\hat{B}}^{\epsilon}_{\epsilon\beta\sigma}$ and $\tilde{\hat{B}}_{\alpha\beta} \stackrel{\text{def.}}{=} \tilde{\hat{B}}^{\epsilon}_{\alpha\beta\epsilon}$.

Table 2: Second Order contracted curvature Tensors

Skew Tensors	Symmetric Tensors
$\tilde{\hat{K}}_{\beta\sigma} \stackrel{\text{def.}}{=} \hat{\varepsilon}_{\sigma\beta} + \hat{\eta}_{\sigma\beta}$ $\overline{\hat{K}}_{\beta\sigma} \stackrel{\text{def.}}{=} \frac{1}{2}(\hat{\varepsilon}_{\sigma\beta} + \hat{\eta}_{\sigma\beta})$ $\hat{K}_{\beta\sigma} \equiv 0$	
$\tilde{\hat{B}}_{[\alpha\beta]} \stackrel{\text{def.}}{=} \hat{\chi}_{\alpha\beta} - \hat{\eta}_{\alpha\beta} - \frac{1}{2}\hat{\varepsilon}_{\alpha\beta}$	$\tilde{\hat{B}}_{(\alpha\beta)} \stackrel{\text{def.}}{=} -\frac{1}{2}\hat{\theta}_{\alpha\beta}$
$\overline{\hat{B}}_{[\alpha\beta]} \stackrel{\text{def.}}{=} \frac{1}{2}\hat{\chi}_{\alpha\beta} - \frac{1}{4}(\hat{\eta}_{\alpha\beta} + \hat{\varepsilon}_{\alpha\beta})$	$\overline{\hat{B}}_{(\alpha\beta)} \stackrel{\text{def.}}{=} -\frac{1}{4}(\hat{\theta}_{\alpha\beta} - \hat{\sigma}_{\alpha\beta} - \hat{\omega}_{\alpha\beta} - \hat{\varpi}_{\alpha\beta})$
$\hat{R}_{[\alpha\beta]} \stackrel{\text{def.}}{=} \frac{1}{2}(\hat{\chi}_{\alpha\beta} - \hat{\eta}_{\alpha\beta} - \hat{\varepsilon}_{\alpha\beta})$	$\hat{R}_{(\alpha\beta)} \stackrel{\text{def.}}{=} \frac{1}{2}(\hat{\psi}_{\alpha\beta} - \hat{\theta}_{\alpha\beta} - \hat{\phi}_{\alpha\beta}) + \hat{\omega}_{\alpha\beta}$

It obvious that, the scalars of the contracted curvature tensors are made of the scalars of the second order tensors (Sec. 3).

Identities

For d-connections $\hat{D}$, $\tilde{D}$, $\overline{D}$ and D , the first and second Bianchi identities are, respectively

$$\begin{aligned}
\text{First, } \quad & \mathfrak{S}_{\mu,\nu,\sigma}(\hat{\Lambda}_{\mu\nu||\sigma}^{\alpha} + \hat{\Lambda}_{\mu\epsilon}^{\alpha}\hat{\Lambda}_{\nu\sigma}^{\epsilon} + \hat{C}_{\mu\epsilon}^{\alpha}N_{\nu\sigma}^{\epsilon}) \equiv 0. \\
& \mathfrak{S}_{\mu,\nu,\sigma}(\hat{\Lambda}_{\nu\mu||\sigma}^{\alpha} + \hat{\Lambda}_{\mu\epsilon}^{\alpha}\hat{\Lambda}_{\nu\sigma}^{\epsilon} + \hat{C}_{\mu\epsilon}^{\alpha}N_{\nu\sigma}^{\epsilon}) \equiv \mathfrak{S}_{\mu,\nu,\sigma}\tilde{\hat{B}}_{\mu\sigma\nu}^{\alpha}. \\
& \mathfrak{S}_{\mu,\nu,\sigma}\hat{C}_{\mu\epsilon}^{\alpha}N_{\nu\sigma}^{\epsilon} \equiv \mathfrak{S}_{\mu,\nu,\sigma}\tilde{\hat{B}}_{\mu\sigma\nu}^{\alpha}. \\
& \mathfrak{S}_{\mu,\nu,\sigma}\hat{C}_{\mu\epsilon}^{\alpha}N_{\nu\sigma}^{\epsilon} \equiv \mathfrak{S}_{\mu,\nu,\sigma}\hat{R}_{\mu\sigma\nu}^{\alpha}.
\end{aligned}$$

$$\begin{aligned}
\text{Second, } \quad 0 & \equiv \mathfrak{S}_{\beta,\nu,\sigma}N_{\nu\sigma}^{\epsilon}(\hat{\Gamma}_{\mu\beta;\epsilon}^{\alpha} - \hat{C}_{\mu\epsilon||\beta}^{\alpha} + \hat{C}_{\mu\rho}^{\alpha}[N_{\beta;\epsilon}^{\rho} - \hat{\Gamma}_{\epsilon\beta}^{\rho}]) \\
& \mathfrak{S}_{\beta,\nu,\sigma}\tilde{\hat{B}}_{\mu\nu\sigma||\beta}^{\alpha} \equiv \mathfrak{S}_{\beta,\nu,\sigma}\{\tilde{\hat{B}}_{\mu\epsilon\sigma}^{\alpha}\hat{\Lambda}_{\nu\beta}^{\epsilon} + N_{\nu\sigma}^{\epsilon}(\tilde{\hat{\Gamma}}_{\mu\beta;\epsilon}^{\alpha} - \hat{C}_{\mu\epsilon||\beta}^{\alpha} + \hat{C}_{\mu\rho}^{\alpha}[N_{\beta;\epsilon}^{\rho} - \tilde{\hat{\Gamma}}_{\epsilon\beta}^{\rho}])\} \\
& \mathfrak{S}_{\beta,\nu,\sigma}\tilde{\hat{B}}_{\mu\nu\sigma||\beta}^{\alpha} \equiv \mathfrak{S}_{\beta,\nu,\sigma}N_{\nu\sigma}^{\epsilon}(\hat{\Gamma}_{(\mu\beta);\epsilon}^{\alpha} - \hat{C}_{\mu\epsilon||\beta}^{\alpha} + \hat{C}_{\mu\rho}^{\alpha}[N_{\beta;\epsilon}^{\rho} - \hat{\Gamma}_{(\epsilon\beta)}^{\rho}]) \\
& \mathfrak{S}_{\beta,\nu,\sigma}\hat{R}_{\mu\nu\sigma;\beta}^{\alpha} \equiv \mathfrak{S}_{\beta,\nu,\sigma}N_{\nu\sigma}^{\epsilon}(\{\hat{\alpha}_{\mu\beta}^{\epsilon}\}_{;\epsilon} - \hat{C}_{\mu\epsilon;\beta}^{\alpha} + \hat{C}_{\mu\rho}^{\alpha}[N_{\beta;\epsilon}^{\rho} - \{\hat{\rho}_{\epsilon\beta}^{\rho}\}])
\end{aligned}$$

5 The W-Tensor

In constructing field theories, a variational method is usually applied to a Lagrangian function having certain properties. In the case of geometric field theories, especially for gravity theories, the Lagrangian usually used is the scalar curvature R , or a general function $f(R)$ depending mainly on the curvature tensor (cf. Ref. 17). In the case of geometries with non-vanishing torsion (T) some authors (cf. Ref. 18, 19) attempted to explore the role of using $f(T)$, in place of $f(R)$, seeking a solution for problems facing gravity theories. In our point of view, it is better to use a Lagrangian, function of both curvature and torsion, as each object plays a role, different from that of the other, with possible mutual interaction.

In most known geometries, curvature and torsion are two distinct entities distinguished by two different tensors. An important object that reflects both curvature and torsion properties is the W-tensor [20]. It can be considered as an amalgam made of both curvature and torsion. This tensor has been first defined [21] and used to construct a unified field theory [22]. It can be constructed, only, in an AP-structure or one of its modifications. In what follows, we are going to construct different forms of this tensor, in the FAP-space in order to make the geometry ready for applications. Let us write the following commutation relations, corresponding to different H- and V- derivatives,

$$\hat{\lambda}_{\mu||\alpha\beta} - \hat{\lambda}_{\mu||\beta\alpha} = \hat{\lambda}_{\epsilon}^{\alpha}\hat{W}_{\mu\alpha\beta}^{\epsilon}, \quad (50)$$

$$\hat{\lambda}_{\mu||\alpha\beta} - \hat{\lambda}_{\mu||\beta\alpha} = \hat{\lambda}_{\epsilon}^{\alpha}\tilde{\hat{W}}_{\mu\alpha\beta}^{\epsilon}, \quad (51)$$

$$\hat{\lambda}_{\mu||\alpha\beta} - \hat{\lambda}_{\mu||\beta\alpha} = \hat{\lambda}_{\epsilon}^{\alpha}\overline{\hat{W}}_{\mu\alpha\beta}^{\epsilon}, \quad (52)$$

$$\hat{\lambda}_{\mu;\alpha\beta} - \hat{\lambda}_{\mu;\beta\alpha} = \hat{\lambda}_{\epsilon}^{\alpha}\overset{o}{\hat{W}}_{\mu\alpha\beta}^{\epsilon}, \quad (53)$$

$$\hat{\lambda}_{\mu|\alpha\beta} - \hat{\lambda}_{\mu|\beta\alpha} = \hat{\lambda}_{\epsilon}^{\alpha}\overset{\diamond}{\hat{W}}_{\mu\alpha\beta}^{\epsilon}, \quad (54)$$

where, when multiplying both sides of the above equations by $\hat{\lambda}_i^\sigma$, we obtain the following definitions of the W-tensor in the FAP-structure, respectively,

$$\hat{W}^{\sigma}_{\cdot\mu\alpha\beta} \stackrel{\text{def.}}{=} \hat{\lambda}_i^\sigma (\hat{\lambda}_{i+}^{\mu||\alpha\beta} - \hat{\lambda}_{i+}^{\mu||\beta\alpha}) \quad (55)$$

$$\widetilde{\hat{W}}^{\sigma}_{\cdot\mu\alpha\beta} \stackrel{\text{def.}}{=} \hat{\lambda}_i^\sigma (\hat{\lambda}_{i-}^{\mu||\alpha\beta} - \hat{\lambda}_{i-}^{\mu||\beta\alpha}) \quad (56)$$

$$\overline{\hat{W}}^{\sigma}_{\cdot\mu\alpha\beta} \stackrel{\text{def.}}{=} \hat{\lambda}_i^\sigma (\hat{\lambda}_{i-}^{\mu||\alpha\beta} - \hat{\lambda}_{i-}^{\mu||\beta\alpha}) \quad (57)$$

$$\overset{o}{\hat{W}}^{\sigma}_{\cdot\mu\alpha\beta} \stackrel{\text{def.}}{=} \hat{\lambda}_i^\sigma (\hat{\lambda}_{i\hat{\mu}\hat{\alpha}\hat{\beta}} - \hat{\lambda}_{i\hat{\mu}\hat{\beta}\hat{\alpha}}) \quad (58)$$

$$\hat{\hat{W}}^{\sigma}_{\cdot\mu\alpha\beta} \stackrel{\text{def.}}{=} \hat{\lambda}_i^\sigma (\hat{\lambda}_{i-}^{\mu||\alpha\beta} - \hat{\lambda}_{i-}^{\mu||\beta\alpha}) \quad (59)$$

Theorem 4. *The curvature and W-tensors have the following coincidences:*

(i) $\overline{\hat{W}}^{\sigma}_{\cdot\mu\alpha\beta}$ coincides with the curvature tensor $\overline{\hat{B}}^{\sigma}_{\cdot\mu\alpha\beta}$.

(ii) $\overset{o}{\hat{W}}^{\sigma}_{\cdot\mu\alpha\beta}$ coincides with the curvature tensor $\hat{R}^{\sigma}_{\cdot\mu\alpha\beta}$.

Proof.

(i) Using (15), we can write (57) in the form

$$\overline{\hat{W}}^{\sigma}_{\cdot\mu\alpha\beta} \stackrel{\text{def.}}{=} \frac{1}{2}(\hat{\Lambda}^{\sigma}_{\cdot\mu\alpha||\beta} - \hat{\Lambda}^{\sigma}_{\cdot\mu\beta||\alpha}) + \frac{1}{4}(\hat{\Lambda}^{\nu}_{\cdot\mu\beta}\hat{\Lambda}^{\sigma}_{\cdot\alpha\nu} - \hat{\Lambda}^{\nu}_{\cdot\mu\alpha}\hat{\Lambda}^{\sigma}_{\cdot\beta\nu}), \quad (60)$$

since it can easily shown that,

$$\hat{\Lambda}^{\alpha}_{\cdot\mu\nu||\sigma} = \hat{\Lambda}^{\alpha}_{\cdot\mu\nu||\sigma} + \frac{1}{2}\mathfrak{S}_{\mu,\nu,\sigma}\hat{\Lambda}^{\epsilon}_{\cdot\mu\nu}\hat{\Lambda}^{\alpha}_{\cdot\sigma\epsilon}.$$

Then substituting into (60), (i) follows.

(ii) Using (28), we can write (58) in the form,

$$\overset{o}{\hat{W}}^{\sigma}_{\cdot\mu\alpha\beta} \stackrel{\text{def.}}{=} \hat{\gamma}^{\sigma}_{\cdot\mu\alpha||\beta} - \hat{\gamma}^{\sigma}_{\cdot\mu\beta||\alpha} + \hat{\gamma}^{\nu}_{\cdot\mu\beta}\hat{\gamma}^{\sigma}_{\cdot\nu\alpha} - \hat{\gamma}^{\nu}_{\cdot\mu\alpha}\hat{\gamma}^{\sigma}_{\cdot\nu\beta} + \hat{\gamma}^{\sigma}_{\cdot\mu\nu}\hat{\Lambda}^{\nu}_{\cdot\alpha\beta}, \quad (61)$$

but we have the relation,

$$\hat{\gamma}^{\alpha}_{\cdot\mu\nu||\sigma} = \hat{\gamma}^{\alpha}_{\cdot\mu\nu||\sigma} + \hat{\gamma}^{\alpha}_{\cdot\mu\epsilon}\hat{\Lambda}^{\epsilon}_{\cdot\sigma\nu},$$

which by substitution into (61) proves the second part of the theorem.

As a consequence of identities (17) and (18), we get $\hat{W}^{\sigma}_{\cdot\mu\alpha\beta} \equiv 0$ and $\hat{\hat{W}}^{\sigma}_{\cdot\mu\alpha\beta} \equiv 0$.

6 Discussion and Concluding Remarks

- (1) The structure given in this article shows that all geometric objects, including the non-linear connection(4), are define from the building blocks of the FAP-structure i.e. the vector fields $\hat{\lambda}_\mu$.
- (2) The torsion corresponding to the non-linear connection $(N^\alpha_{\beta;\sigma} - N^\alpha_{\sigma;\beta})$ coincides with the torsion of the Bervald type linear connection $\hat{T}^\alpha_{\mu\nu}$ given by (31).
- (3) The anti-curvature tensor defined in an AP-structure has been shown to be useful in describing antigravity [16]. It is made of the torsion of the AP-structure. The definition of this tensor, in the FAP-structure, shows that it contains, implicitly, the non-linear connection. Applications, using FAP will illuminate the role of the non-linear connection in this respect, if any.
- (4) Its W-tensor (Sec. 5) reflects, not only the effects of the curvature and torsion (as in the case of AP-structure [20]), but also the effect of the non-linear connection. This is an important property which enables us to explore the role of non-linear connections, in unifying fundamental interactions, if any.
- (5) The following table (Table 3) gives a brief comparison between the FAP-, the AP-, Finsler and Riemannian structures.

Table 3: Comparison between FAP-, AP-, Finsler and Riemann geometries

Criteria	FAP	AP	Finsler	Riemann
Underlying Space	(M, L_i)	(M, λ_i)	(M, F)	(M, g)
Structure Dependence	$L_i(x, y)$	$\lambda_i(x)$	$F(x, y)$	$g(x)$
Number of Natural Metric Connections	2	2	1	1
Number of built-in Non-Linear Connections	2	0	1	0
Number of Anti-curvature Tensors	1	1	0	0
Number of W-Tensors*	1	1	0	0

*Non-Vanishing and differs from the Curvature tensor.

From this table, it is clear that the FAP-structure is more thorough than both conventional AP and Finsler structures, in particular:

- (i) It is clear that the only non-vanishing W-tensor and different from the curvature tensors is that given by (56). Using (14) this tensor can be written in the form

$$\widetilde{W}^{\sigma}_{\cdot\mu\alpha\beta} \stackrel{\text{def.}}{=} \hat{\Lambda}^{\sigma}_{\cdot\mu\alpha||\beta} - \hat{\Lambda}^{\sigma}_{\cdot\mu\beta||\alpha} + \hat{\Lambda}^{\nu}_{\cdot\mu\beta} \hat{\Lambda}^{\sigma}_{\cdot\alpha\nu} - \hat{\Lambda}^{\nu}_{\cdot\mu\alpha} \hat{\Lambda}^{\sigma}_{\cdot\beta\nu}. \quad (62)$$

Due to the importance of this tensor in physical applications, we are going to write its possible contractions which will be written, in terms of the tensors of Table 1, as

$$\widetilde{S}_{\alpha\beta} \stackrel{\text{def.}}{=} \widetilde{W}^{\epsilon}_{\cdot\epsilon\alpha\beta} = \hat{\epsilon}_{\beta\alpha} + 2\hat{\eta}_{\beta\alpha}, \quad (63)$$

$$\widetilde{W}_{\cdot\mu\alpha} \stackrel{\text{def.}}{=} \widetilde{W}^{\epsilon}_{\cdot\mu\alpha\epsilon} = \hat{\chi}_{\mu\alpha} - \hat{\eta}_{\mu\alpha} - \hat{c}_{\mu||\alpha} - \hat{\varpi}_{\alpha\mu} + \hat{\sigma}_{\alpha\mu} + \hat{\omega}_{\alpha\mu} \quad (64)$$

The tensor (63) is anti-symmetric. So, it cannot be further contracted. The tensor (64) is non-symmetric. So, its symmetric part can be contracted to give

the scalar

$$\widetilde{W} = -\frac{1}{2}\hat{\theta} + 3\hat{\omega} + \hat{\sigma} \quad (65)$$

- (ii) We stress again that, the W-tensor, defined in Sec. 5 contains the effects of the curvature tensor, the torsion of the linear connection, and the non-linear connection. It is to be considered that this tensor has only one value given by (56). Other values (55), (57), (58) are either vanishing (due to (17) and (18)) or coincides with the curvature tensors (Theorem 4). This is an important property, of the FAP-structure, which would lead to the construction of a unique field theory using this tensor.

The following schematic diagram shows the relations between the above mentioned geometric structures. Solid arrows represent the direction towards special cases, while dashed arrows show the associated structures.




Fig. 1: Relation between different geometric structures.

References

- [1] I. Bucataru and R. Miron, *Finsler-Lagrange Geometry. Applications to Dynamical Systems*, Editure of Romanian Academy, Bucharest (2007).
- [2] S.I. Vacaru, *Int.J.Geom.Meth.Mod.Phys.* **05**, 473 (2008); arXiv:0801.4958v1.
- [3] X. Li and Z. Chang, *Phys.Lett.B* **692**, 1 (2010); arXiv:0909.3713.
- [4] M.I.Wanas, *Stud.Cercet.Stiin.Ser.Mat.Univ.Bacau* No.10, 297 (2001); arXiv:gr-qc/0209050.
- [5] J.F. da Rocha-Neto, J.W. Maluf and S. C. Ulhoa, *Phys.Rev.D* **82**, 124035 (2010); arXiv:1101.2425.
- [6] A.A. Sousa, R.B. Pereira and A.C. Silva, *Grav.Cosmol.* **16**, 25 (2010); arXiv:0803.1481.
- [7] S.G. Turyshev and V.T. Toth, *Living Rev. Relativity* **13**, 4 (2010); <http://relativity.livingreviews.org/>.

- [8] E. Battaner and E. Florido, *Fund. Cosmic Phys.* **21**, 1 (2000); arXiv:astro-ph/0010475.
- [9] J.L. Tonry, P.B. Schmid et al., *Astrophys.J.* 594, 1 (2003); arXiv:astro-ph/0305008.
- [10] M.I. Wanas, *Modern Physics Letters A* **24**, No. 22, 1749 (2009); arXiv:0801.1132.
- [11] F.I. Mikhail, *Ain Shams Sci. Bul.* **6**, 87 (1962).
- [12] F.I. Mikhail and M.I. Wanas, *Int. J. Theor. Phys.* **20**, 671 (1981).
- [13] M.I. Wanas, *Int.J.Geom.Meth.Mod.Phys.***4**, 373 (2007); arXiv:gr-qc/0703036.
- [14] R. Miron and M. Anastasiei, *Vector bundles and Lagrange spaces with applications to relativity*, Vol. 1, Geometry Balkan Press, Bucurest, Romania (1997).
- [15] M.I. Wanas, *Astrophys.Space Sci.* **258**, 237 (1998); arXiv:gr-qc/9904019v1.
- [16] M.I. Wanas, Six Conf. "Nuclear and Particle Physics " held in Luxor, Egypt 17-21 Nov. (2007); arXiv:0809.5040.
- [17] A. De Felice and S. Tsujikawa, *Living Rev. Rel.* **13**, 3 (2010); arXiv:1002.4928.
- [18] G.R. Bengochea and R. Ferraro, *Phys. Rev. D* **79**, 124019 (2009); arXiv:0812.1205.
- [19] R. Myrzakulov; arXiv:1006.1120.
- [20] N.L. Youssef and A.M. Sid-Ahmed, *Rept.Math.Phys.* **60**, 39 (2007); arXiv:gr-qc/0604111.
- [21] M.I. Wanas, *A generalized field theory and its applications in cosmology*, Ph. D. Thesis, Cairo University, (1975).
- [22] F.I. Mikhail and M.I. Wanas, *Proc. R. Soc. London, A.* **356**, 471 (1977).